

Clinical Trial	Trial Length	Number of Patients on Risperdal	Reports of Gynecomastia	Gynecomastia Rate
AGES NED-9 (11-18)	10 weeks	19	0	0%
BEL-22 (3-14)	4 weeks	7	0	0%
BEL-24 (6-14)	4 weeks	6	0	0%
INT-41 (4-14)	1 year	504	24	4.8%
USA-93 (5-12)	7 weeks	55	0	0%
USA-97 (5-12)	1 year	107	1	0.9%
CAN-19 (5-12)	7 weeks	53	0	0%
CAN-20 (5-12)	1 year	77	0	0%
USA-150 (5-17)	8 weeks	49	1	2%
CAN-23 (5-12)	8 weeks	41	0	0%
HUN-4 (6-16)	2 years	35	0	0%
INT-70 (6-14)	1 year	48	3	6%
USA-231 (13-17)	8 weeks	279	2	0.7%
USA-234 (13-17)	6 months	297	4	1%
INT-79 (5-17)	9 months	527	8	1.5%
INT-84 (6-18)	1 year	232	2	0.8%
BIM-301 (10-17)	3 weeks	111	0	0%
SCH-302 (13-17)	6 weeks	106	0	0%

Source: 2-25

EXHIBIT
 Date: 17B
 M.L. GRAY, CSR, RPR

PLAINTIFF'S
 EXHIBIT
 17B